

Table 13. Proportion of persons surviving for more than 12, 24, and 36 months after an AIDS diagnosis in 2000, by selected characteristics—United States

| | No. of persons | Survival in months | | | |
|--|----------------|--------------------|-------------|-------------|--|
| | | >12 | >24 | >36 | |
| Proportion | | | | | |
| Age at diagnosis (yrs) | | | | | |
| <13 | 118 | 0.96 | 0.95 | 0.95 | |
| 13–14 | 60 | 1.00 | 0.98 | 0.93 | |
| 15–19 | 289 | 0.98 | 0.95 | 0.93 | |
| 20–24 | 1,302 | 0.95 | 0.92 | 0.89 | |
| 25–29 | 3,367 | 0.94 | 0.91 | 0.89 | |
| 30–34 | 6,314 | 0.94 | 0.90 | 0.88 | |
| 35–39 | 8,676 | 0.92 | 0.88 | 0.86 | |
| 40–44 | 7,245 | 0.90 | 0.86 | 0.83 | |
| 45–49 | 4,959 | 0.88 | 0.83 | 0.79 | |
| 50–54 | 2,833 | 0.86 | 0.80 | 0.76 | |
| 55–59 | 1,438 | 0.84 | 0.79 | 0.75 | |
| 60–64 | 774 | 0.79 | 0.74 | 0.70 | |
| ≥65 | 689 | 0.73 | 0.66 | 0.61 | |
| Race/ethnicity | | | | | |
| White, not Hispanic | 10,994 | 0.91 | 0.87 | 0.85 | |
| Black, not Hispanic | 18,828 | 0.90 | 0.85 | 0.81 | |
| Hispanic | 7,598 | 0.91 | 0.87 | 0.85 | |
| Asian/Pacific Islander | 340 | 0.91 | 0.89 | 0.88 | |
| American Indian/Alaska Native | 168 | 0.92 | 0.89 | 0.85 | |
| Transmission category | | | | | |
| Male adult or adolescent | | | | | |
| Male-to-male sexual contact | 12,777 | 0.92 | 0.89 | 0.87 | |
| Injection drug use | 5,406 | 0.87 | 0.82 | 0.78 | |
| Male-to-male sexual contact and injection drug use | 1,768 | 0.92 | 0.87 | 0.83 | |
| Heterosexual contact | 2,963 | 0.91 | 0.87 | 0.84 | |
| Other ^a | 4,975 | 0.87 | 0.83 | 0.81 | |
| Subtotal | 27,889 | 0.90 | 0.86 | 0.84 | |
| Female adult or adolescent | | | | | |
| Injection drug use | 2,560 | 0.89 | 0.83 | 0.77 | |
| Heterosexual contact | 4,782 | 0.92 | 0.88 | 0.85 | |
| Other ^a | 2,715 | 0.90 | 0.85 | 0.82 | |
| Subtotal | 10,057 | 0.91 | 0.86 | 0.82 | |
| Child (<13 yrs at diagnosis) | | | | | |
| Perinatal | 104 | 0.95 | 0.94 | 0.94 | |
| Other ^b | 14 | 1.00 | 1.00 | 1.00 | |
| Subtotal | 118 | 0.96 | 0.95 | 0.95 | |
| Total^c | 38,064 | 0.90 | 0.86 | 0.83 | |

Note. Excludes persons whose date of death is before, or in the same month as, date of diagnosis. Excludes persons whose month of diagnosis or month of death is unknown.

^a Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

^b Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^c Includes 136 persons of unknown race or multiple races.